



Italian ryegrass

Lolium perenne L. ssp. multiflorum (Lam.) Husnot.

Bagavathiannan, M¹, Maity, A¹, Ackroyd, V², Flessner, M³, Rubione, CG⁴, VanGessel, MJ⁴

¹Texas A&M University

²USDA-ARS, Beltsville, MD

³Virginia Tech

⁴University of Delaware Research & Education Center

Also known as: *annual ryegrass, annual rye.*

Biology

Italian ryegrass is a winter annual weed species found throughout the US infesting small grain production, where it can cause yield losses in excess of 50%, reaching as high as 100% in severely infested fields. In many parts of the south-central US, a second flush of ryegrass seedlings emerges during the spring, which creates difficulties for management. Ryegrass populations have different degrees of seed dormancy and thus show a staggered but continuous emergence pattern until the shade from the cash crop completely suppresses their emergence. Italian ryegrass exhibits high levels of genetic diversity, which allows for adaptation to a wide range of soil textures and environmental conditions. Because it is an annual weed, it must regenerate from seed each year. Seeds can germinate across a wide range of temperatures, but

soil temperatures of approximately 65-70 degrees F are considered optimal for germination in the south-central US. Seedlings show rapid growth potential, with the production of multiple tillers. Italian ryegrass tends to dominate conservation tillage systems where soil disturbance is limited.

Italian ryegrass often displays high levels of seed shattering. Research conducted at Washington State University found that only 58% of weed seeds are retained at wheat harvest. Results from Texas also confirm these findings. However, the seed retained on the plants at harvest can still be substantial. In Virginia, Italian ryegrass has been documented to retain 110 to 293 lb. of seed per acre at wheat harvest, well exceeding recommended seeding rates of about 25 lb. per acre for the species when used as a forage. Seedbank persistence is not well researched, but forage systems requiring annual re-seeding suggest that seeds are relatively short-lived.

Italian ryegrass is a wind-pollinated species, capable of dispersing pollen over long distances. Though studies are limited, it is likely that pollen- dispersal can spread herbicide resistance across ryegrass populations in an agricultural landscape.



Winter wheat heavily infested with Italian ryegrass, Texas. Photo credit: Claudio Rubione, University of Delaware.


How is the lifecycle of a italian ryegrass plant?

Seeds start germinating in fall followed by a rapid growth rate when the temperature is moderately low. During winter, the growth slows down and reaccelerates in spring. Flowering occurs during May to July depending on the climatic conditions and produces a large quantity of seeds which shatter substantially during maturity. Usually the plants die after seed maturity and do not develop a sizable seedbank due to short seed dormancy period, which ensures its resurgence in fall.

Italian ryegrass lifecycle



Herbicide resistance



This graph shows reported herbicide-resistance for giant ragweed in the US; both resistance to a single herbicide group as well as resistance to multiple herbicide groups. Graph is based on reports to the International Herbicide-Resistant Database, www.weedscience.org and published research. Contact your local extension office for details about resistance in your area and management options.

*Herbicide names listed are representative products that contain specific active ingredients. Last updated on: 7-10-2020

Integrated weed management strategies



Click here to watch the video: <https://www.youtube.com/watch?v=Gyv0zbKpFfU>

Cultural such as [crop rotation](#) and [narrow row spacing](#) can reduce Italian ryegrass populations. Crop rotation out of continuous wheat will shift typical production practices like the timing of planting,

tillage, and herbicide applications and allow for more diverse management tactics. Italian ryegrass is intolerant to shade so narrowing row spacing allows crops to canopy quickly and slows down the growth of Italian ryegrass.

Tillage, herbicide application, and [harvest weed seed control](#) are other key tactics to prevent Italian ryegrass from setting seed. [Crop rotation](#) out of continuous wheat, adding summer annual crop with an effective burndown application is a potential alternative management strategy. Till or spray early enough in spring so the Italian ryegrass plants do not set seed.

Cover crops If Italian ryegrass infests your [cover crops](#), make sure the cover crops are terminated before the weeds set seeds. Cover crops are not the best option for suppression of Italian ryegrass compared to other summer annual weeds. If ryegrasses are a weed of concern in your cropping system, consider planting cover crops other than annual ryegrass and buy clean cover crop seed.

Mechanical Pulling, cutting, or disking – the traditional mechanical management methods – do not provide sufficient control since ryegrass is highly tolerant of them. However, small patches can be controlled by hand-weeding, which is not a viable option for large scale production scenarios. Trampling, mowing, or grazing are also equally ineffective since ryegrass can easily regrow with any available soil moisture. Occasional deep tillage may stop ryegrass seed germination by burying them at greater depth, but takes up to seven years to achieve that.

Harvest weed seed control (HWSC) tactics developed in Australia, another country where ryegrass is the major weed problem in wheat, promises great potential to reduce ryegrass seedbank inputs and subsequent field infestations. [HWSC](#) aims to destroy the weed seeds during crop harvest so that the soil seedbank declines rapidly,

gradually drawing down the weed seedbank and making it easier to manage with available herbicide programs. High weed seed retention is a prerequisite for successful HWSC. Currently, three types of HWSC are in use in Australia and are being tested for their potential across the US. The first type, narrow-windrow burning, collects the weed seeds and cash crop chaff from the harvester and deposits it in narrow lines in the field, which are then burnt. The burning is of concern from an environmental pollution and uncontrolled fire risk standpoint. To avoid such problems, chaff lining involves collecting the chaff fraction separately from the straw in a similar way, then the seeds rot or germinate and then die due to poor seed-soil contact in the windrows. The latest development is the impact mill technology, which has been designed to collect the weed seeds separately from the chaff and grind them for complete destruction using high speed impact mills at the back of the combine. See our video, which details a 3-year study conducted in Texas and Arkansas showing that HWSC (narrow-windrow burning), when combined with the best herbicide program existing in that region, achieves maximum ryegrass control and reduces the soil seedbank significantly.

Chemical control remains the most-used tool to combat Italian ryegrass. However, as previously noted, Italian ryegrass resistance has limited the availability of key herbicides.

Chemical control of Italian ryegrass in small grains has often relied on Group 2 herbicides, such as mesosulfuron or pyroxasulfone (Osprey or PowerFlex) and Group 1 (Axial XL, pinoxaden); where resistance is widespread pyroxasulfone (Anthem Flexx or Zidua) is commonly used. Pyroxasulfone needs to be applied prior to Italian ryegrass emergence since it will not control emerged plants.

Though glyphosate resistance has been reported in seven states, it is still used heavily as a burndown herbicide to control Italian ryegrass prior to planting

summer crops. Controlling glyphosate-susceptible Italian ryegrass in the spring prior to planting corn, cotton, or soybean is challenging due to its vigorous growth and potential antagonism when glyphosate is mixed with other herbicides. Paraquat is utilized extensively as a burndown in the southern states for Italian ryegrass.

In broadleaf crops, postemergence grass herbicides (such as clethodim and sethoxydim) can be used as part of a burndown program or as a postemergence application. Most of the Group 15 herbicides (such as Dual Magnum, Zidua) will provide residual control, yet their utility is limited since most Italian ryegrass populations will have emerged at time of summer crop planting.

Refer to your local extension’s weed management guide for effective options.

There are no reliable microbial-based **biological** control options for Italian ryegrass. Though many fungi are reported to infect ryegrass seedheads, they do not sufficiently and consistently reduce seed production enough to represent a management option. However, research conducted in Texas showed that delaying field disking after wheat harvest can increase ryegrass seed loss from the soil surface by exposing them to ants, beetles, birds and other animals that consume them.

Similar weed species

Rigid ryegrass (*L. rigidum*) and perennial ryegrass (*L. perenne*) are the other two closely related species of Italian ryegrass. As the Italian and perennial are the two most commonly occurring species in the States and can cross easily to each other giving rise to a diverse range of phenotypes, here is a comparison table based on a few key identifying traits.



To differentiate Italian ryegrass from other grass species, look closely for its conspicuous claw-like auricles in the collar region of the stem. Picture credit: Claudio Rubione, University of Delaware.

Characteristics	<i>L. perenne</i> or Perennial Type	<i>L. multiflorum</i> or Annual Type
Growth habit	Mostly spreading type with relatively less vigorous growth. Plants have a bunched form, with numerous long, narrow, stiff leaves near the base of the plant. Picture credit: Aniruddha Malty, Texas A&M University	Mostly erect to semi-erect type with highly vigorous growth. Plants look straight and less bunched with wider and fewer leaves. Picture credit: Aniruddha Malty, Texas A&M University
Auricle: it is a small finger-like projection from the base of the leaf where leaf sheath attaches to the stem.	Very short and they do not clasp the stem. Picture credit: North Carolina State University	The auricles are long and clasp around the stem. Picture credit: North Carolina State University
Seed and seed awn: Awn is a hair- or bristle-like appendage on top of the seed.	Smaller seeds, mostly 6 to 10 flowers per spikelet. Awn is usually absent.	Larger seeds, 10 to 20 flowers per spikelet. Awn is usually present. Picture credit: Aniruddha Malty, Texas A&M University

For more information on Italian ryegrass management, visit:

<https://iwilltakeaction.com/weed/italian-ryegrass>



References:

- Galla M, Italian ryegrass management in Sacramento Valley wheat <https://ucanr.edu/blogs/blogcore/postdetail.cfm?postnum=26333> Accessed: August, 2 2021.
- Heap I (2019) The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org/Home.aspx> Accessed: August 29, 2021.
- Hefty B, Hefty D, Weed of the week: Italian ryegrass <http://watch.agphd.com/detail/videos/weed-of-the-week/video/5557846392001/weed-of-the-week-1012-italian-ryegrass-air-date-8-27-17> Accessed: Aug 2, 2021.
- Take action on weeds: Italian <https://iwilltakeaction.com/weed/italian-ryegrass> Accessed August 2, 2021.
- Beam SC, Mirsky SB, Cahoon C, Haak D (2019) Harvest weed seed control of Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot], common ragweed (*Ambrosia artemisiifolia* L.), and Palmer amaranth (*Amaranthus palmeri* S. Watson), Weed Technology: 24 doi.org/10.1017/wet.2019.46
- Bararpour MT, Norsworthy JK, Burgos NR, Korres NE, Gbur EE (2019) Identification and biological characteristics of ryegrass (*Lolium* spp.) accessions in Arkansas. Weed Technology 33:627-632 doi.org/10.1017/wsc.2016.28
- Brunnharo CA, Bradley DH, (2018) Multiple herbicide-resistant Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot] in California perennial crops: characterization, mechanism of resistance, and chemical management. Weed Science 66:696-701 doi.org/10.1017/wsc.2018.50
- Liu M, Hulting A, Mallory-Smith C (2017) Characterization of multiple herbicide-resistant Italian ryegrass (*Lolium perenne* ssp. *multiflorum*) populations from winter wheat fields in Oregon. Weed Science 64:331-338 doi.org/10.1614/WS-D-15-00147.1
- Lyon DJ, Huggins DR, Spring JF (2016) Windrow burning eliminates Italian ryegrass (*Lolium perenne* ssp. *multiflorum*) seed viability. Weed Technology 30: 279-283 doi.org/10.1614/WT-D-15-00118.1
- Mahajan G, McKenzie K, Chauhan BS (2019) Influence of row spacing and cultivar selection on annual ryegrass (*Lolium rigidum*) control and grain yield in chickpea (*Cicer arietinum*). Crop & Pasture Science 70:140-146 doi.org/10.1071/CP18436
- Walsh MJ, Broster JC, Schwartz-Lazaro LM, Norsworthy JK, Davis AS, Tidemann BD, Beckie HJ, Lyon DJ, Soni N, Neve P, Bagavathiannan MV (2018) Opportunities and challenges for harvest weed seed control in global cropping systems. Pest Management Science 74:2235-2245 doi.org/10.1002/ps.4802
- Cite as: Bagavathiannan, M., Maity, A., Ackroyd, V., Flessner, M., Rubione, C.G., VanGessel, M.J. (2021). Italian Ryegrass. Getting Rid of Weeds Through Integrated Weed Management. www.growiwm.org



Funding for this project is from USDA Area-Wide Management of Agricultural Pests.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident. USDA is an equal opportunity provider, employer, and lender.